

FULL WEIGHT-BEARING AFTER DUAL-FIXATION OF CLAVICLE FRACTURES APPEARS SAFE AND EFFECTIVE: A MULTI-CENTER COMPARATIVE STUDY

Jack Twomey-Kozak, BS*¹; Nicholas Morris, BA¹; Keith G. Whitlock, MD¹; Christian A. Pean, MD; Julius A. Bishop, MD³; Michael J. Gardner MD³; Mark J Gage, MD^{2,4}; Oke A. Anakwenze, MD, MBA, ¹Malcolm DeBaun, MD².

¹Department of Orthopaedic Surgery, Duke University Medical Center, Durham, NC. ²Division of Orthopaedic Trauma, Department of Orthopaedic Surgery, Duke University Medical Center, Durham, NC, USA. ³Department of Orthopaedic Surgery, Stanford University School of Medicine, Stanford, CA, United States. ⁴R Adams Cowley Shock Trauma Center, University of Maryland School of Medicine, Baltimore, MD

INTRODUCTION

Recently, dual-plating of displaced clavicle fractures has become a popular construct to reduce implant prominence while optimizing balanced fixation. Traditionally patients have a limited weight bearing period after clavicle fracture fixation. However, dual-plate fixation may allow for immediate weight bearing and facilitate earlier rehabilitation, especially in polytraumatized patients. The current study aimed to evaluate healing and complication rates between different weight bearing protocols following dual-plating of displaced diaphyseal clavicle fractures.

METHODS


There were 69 patients from two separate level 1 trauma centers who sustained a diaphyseal clavicle fracture and were subsequently treated with dual-plate fixation from 2014 to 2022. Patients were included with minimum of one year follow up or until radiographic and clinical union. Patients were either non-weight-bearing (NWB) (n = 11), Partial-weight-bearing (PWB) (n = 36), or weight-bearing-as-tolerated (WBAT) (n = 22) based on surgeon preference. Complications, re-operation rates secondary to loss of reduction/fixation, and union rates were compared between weight-bearing groups using Fisher's exact test.

RESULTS

There was no significant difference in union rates (NWB 100%, PWB 97.2%, WBAT 95.45%). There was no significant difference in overall re-operation rate between weight-bearing groups. The majority of reoperations were due to symptomatic hardware removal. One patient in the PWB group required revision surgery for explicit nonunion. One patient in the WBAT group required I&D for superficial wound infection.

DISCUSSION/CONCLUSION

Patients treated with dual-plate fixation had well-balanced and stable fixation constructs. Full weight bearing after dual-plate fixation for displaced diaphyseal clavicle fractures may be safe and effective. In comparison, limited weight bearing does not seem to offer any clinical benefit related to union and complication rates. Allowing patients to weight bear immediately after clavicle fracture fixation may improve rehabilitation especially in polytraumatized patients who require crutch/walker weight bearing for concomitant injuries.



*Note: Reoperation (1B) etiology includes loss of reduction/fixation, removal of symptomatic hardware, infection, heterotopic ossification, or nonunion requiring revision