

2.4. DIAGNOSIS: PATHOGEN ISOLATION

Authors: Fernando Motta, William Li

QUESTION 1: Is there a method to detect sessile microorganisms that have resulted in an infection following orthopaedic procedures?

RECOMMENDATION: Yes. Molecular techniques such as polymerase chain reaction (PCR), next-generation sequencing (NGS) and synovial biomarkers such as alpha-defensin or leukocyte esterase have been shown to be powerful tools in detecting prosthetic joint infections (PJI) with negative cultures, although conflicting data exists on PCR. Sonication of explanted prosthetics can enhance both the sensitivity of conventional cultures and PCR.

LEVEL OF EVIDENCE: Strong

DELEGATE VOTE: Agree:85%, Disagree: 9%, Abstain: 6% (Super Majority, Strong Consensus)

RATIONALE

The colonization of prostheses by sessile bacteria is a feared complication of orthopaedic procedures. These microorganisms anchor themselves to the surface of prosthetic implants and form a colony of immobile bacteria cross-linked by an extracellular matrix of polymeric substances, known as biofilm [1]. The presence of biofilm on prosthetic implants, especially that of prosthetic joints, makes both detection and treatment of infections difficult [2]. While there is no gold standard for definitive diagnosis of PJI, a multi-criteria definition created by Musculoskeletal Infection Society (MSIS) is often used to diagnose PJI [3,4]. The MSIS criteria utilizes the obtaining of cultures of joint aspirate or periprosthetic tissue as one of the major criteria to prove the presence of pathogens in the prosthetic joint. Unfortunately, cultures can be unreliable when detecting biofilms [5,6]. Intraoperative cultures alone also can have a high rate of contamination and false positives [7]. Thus, alternative methods of confirming the presence of organisms in PJI have been proposed [8,9]. Some of these diagnostic techniques include PCR, NGS, prosthesis sonication and joint biomarkers.

Polymerase Chain Reaction

The use of PCRs to detect bacterial nucleic acids in prosthesis infections can be an effective way of detecting sessile microorganisms otherwise not picked up in cultures [10,11]. PCR sequencing of bacterial ribosomal nucleic acids has shown to have higher sensitivity in detecting bacteria than culture, as well as identifying polymicrobial infections that may not be picked up by culture [12–15]. Jahoda et al. showed that the use of PCR can detect as few as 590 colony forming units of *S. aureus*, making detection of PJI even in the presence of antibiotics feasible [11]. PCR has also shown benefit in detecting genes responsible for biofilm production and methicillin resistance [11,16].

In spite of the literature describing the merits of PCR, there is data suggesting that the efficacy of PCR is not as high as once thought. Studies have suggested that PCR has similar or less sensitivity for detecting bacteria in PJI as traditional cultures [17–20]. PCR has also been shown to have questionable sensitivity over the last years. A meta-analysis performed by Jun et al. looking at online databases from 2013 to 2017 showed that there has been a decrease in pooled sensitivity compared to a previous meta-analysis performed by Qu et al. in 2013 (0.76, (95% confidence interval (CI) 0.65-0.85) vs.0.86, (95% CI 0.77-0.92) respectively), with no change in specificity [21,22].

Next-Generation Sequencing

Recently, NGS has proven to be efficacious in diagnosis of culture-negative PJIs as well. A prospective study performed by Tarabichi et al. evaluated the accuracy of NGS in identifying PJIs in 78 patients undergoing revision or primary arthroplasties. NGS identified infections in 25 of the 28 cases considered to be PJIs by MSIS criteria (95% CI 71.8% to 97.7%), whereas cultures were only able to identify 17 cases (95% CI 40.6% to 78.5%). In cases where both cultures and NGS were positive, NGS showed a high degree of concordance to traditional cultures as well [23].

NGS has also shown high degrees of detection in synovial fluid samples. Another study conducted by Tarabichi et al. analyzed 86 samples of synovial fluid from the hip or knees of patients undergoing PJI evaluation. They found that NGS had a positive result in 10 samples that were culture-negative. Five of these samples had elevated inflammatory biomarkers, indicating an infectious process, while the other five had negative inflammatory biomarkers. These results suggest that NGS may be a valuable tool for evaluating for PJIs in the preoperative setting, but may also be at risk for false positives [24].

In addition to diagnosing prosthetic infections, NGS may also be useful for identification of causative organisms in culture-negative PJIs [23]. Furthermore, the speed at which NGS can explore an entire genome makes it a superior alternative to PCR [25]. While NGS has exciting potential as a powerful diagnostic tool for culture-negative PJIs, there has been limited data showing its effectiveness in diagnosing other prosthetic infections. In addition, there has been no direct comparison between the effectiveness PCR and NGS. Finally, it is important to consider that the high sensitivity may predispose NGS to a high false-positive rate and false diagnosis of PJIs [25].

Sonication

The use of sonication to break up biofilm in prosthetic implants has been shown to increase the sensitivity of both cultures and PCR when testing for infection. A prospective study performed by Tani et al. compared the sensitivity and specificity of cultures obtained from sonicated explants to conventional cultures of periprosthetic tissue in 114 patients who underwent hip and knee revisions due to PJI and aseptic loosening. Sonicated cultures had a significantly-increased sensitivity when compared to conventional cultures (77.0% vs. 55.7%). There were no significant differences in specificity of either detection method [26].

There are some studies suggesting that sonication of prosthesis may improve the diagnosing capacity of PCR in the diagnosis of culture-negative PJIs [27–29]. However, their statistical significance remains controversial. A recent meta-analysis of nine studies looking at the efficacy of sonication in PCR was performed by Liu et al. [30] found that PCR for sonication prosthetic fluid was to have clinically acceptable diagnostic values for detecting PJIs, with a pooled sensitivity of 75% (95% CI 0.71 to 0.79) and specificity of 96% (95% CI 0.94 to 0.97) [30].

Joint Biomarkers

Inflammatory biomarkers in the blood such as erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP), as well as synovial fluid leukocyte esterase have been part of the 2011 MSIS criteria and the 2013 consensus group modification criteria in the diagnosis of PJI [3,31]. The updated MSIS criteria put forth by Parvizi et al. in 2018 added the presence of synovial alpha-defensin and synovial CRP as criteria for diagnosis of PJI [4]. Synovial biomarkers such as leukocyte esterase and alpha-defensin have been shown to have high sensitivity and specificity in diagnosis of PJI, and are more specific than serum inflammatory biomarkers [32–34]. The benefit of these biomarkers are that they are faster and less invasive than traditional cultures. Biomarker assays also do not require tissue sampling and may be performed on synovial fluids, which increases the convenience of these tests in diagnosing PJIs in the preoperative setting. The major drawback of joint biomarkers is that they can only indicate the presence of infection and not its specific nature. Therefore, biomarkers are best utilized as a preliminary indicator of the presence or absence of joint infection. They are best followed up by diagnostic assays such as PCR, NGS or cultures to better determine the nature of infection.

Conclusion

There are a number of methods to detect sessile microorganisms in infections following orthopaedic procedures. The use of PCR in the diagnosis of culture-negative PJI has shown to be more sensitive than traditional cultures but there is conflicting data. The use of inflammatory biomarkers in both the blood in synovial fluid is also effective, but cannot characterize the nature of infection or organism involved. NGS is a new test can determine the presence of sessile microorganisms with more precision and speed than traditional cultures. Finally, sonication of explants has shown to improve the sensitivity of both cultures and PCR in diagnosing prosthesis infections.

REFERENCES

- [1] Costerton JW, Montanaro L, Arciola CR. Biofilm in implant infections: its production and regulation. *Int J Artif Organs*. 2005;28:1062–1068. doi:10.1177/039139880502801103.
- [2] Jacqueline C, Caillon J. Impact of bacterial biofilm on the treatment of prosthetic joint infections. *J Antimicrob Chemother*. 2014;69 Suppl 1:i37–i40. doi:10.1093/jac/dku254.
- [3] Parvizi J, Zmistowski B, Barbari EF, Bauer TW, Springer BD, Della Valle CJ, et al. New definition for periprosthetic joint infection: from the workgroup of the musculoskeletal infection society. *Clin Orthop Relat Res*. 2011;469:2992–2994. doi:10.1007/s11999-011-2102-9.
- [4] Parvizi J, Tan TL, Goswami K, Higuera C, Della Valle C, Chen AF, et al. The 2018 definition of periprosthetic hip and knee infection: The evidence-based and validated criteria. *J Arthroplasty*. 2018;33:1309–1314. doi:10.1016/j.arth.2018.02.078.
- [5] McConoughey SJ, Howlin R, Granger JF, Manning MM, Calhoun JH, Shirtliff M, et al. Biofilms in periprosthetic orthopedic infections. *Future Microbiol*. 2014;9:987–1007. doi:10.2217/fmb.14.64.
- [6] Neut D, Van Horn JR, Van Kooten TG, Van Der Mei HC, Busscher HJ. Detection of biomaterial-associated infections in orthopaedic joint implants. *Clin Orthop Relat Res*. 2003;261–268. doi:10.1097/01.blo.0000073345.50837.84.
- [7] Barrack RL, Aggarwal A, Burnett RS, Clohisy JC, Ghanem E, Sharkey P, et al. The fate of the unexpected positive intraoperative cultures after revision total knee arthroplasty. *J Arthroplasty*. 2007;22:94–99. doi:10.1016/j.arth.2007.03.029.
- [8] Gbejuade HO, Lovering AM, Webb JC. The role of microbial biofilms in prosthetic joint infections. *Acta Orthop*. 2015;86:147–158. doi:10.3109/17453674.2014.966290.
- [9] Patel R, Alijanipour P, Parvizi J. Advancements in diagnosing periprosthetic joint infections after total hip and knee arthroplasty. *Open Orthop J*. 2016;10:654–661. doi:10.2174/1874325001610010654.
- [10] Bergin PF, Doppelt JD, Hamilton WG, Mirick GE, Jones AE, Sritulanondha S, et al. Detection of periprosthetic infections with use of ribosomal RNA-based polymerase chain reaction. *J Bone Joint Sur Am*. 2010;92:654–663. doi:10.2106/JBJS.I.00400.
- [11] Jahoda D, Landor I, Benedík J, Pokorný D, Judl T, Barták V, et al. PCR diagnostic system in the treatment of prosthetic joint infections. *Folia Microbiol (Praha)*. 2015;60:385–391. doi:10.1007/s12223-014-0370-y.
- [12] Suda AJ, Kommerell M, Geiss HK, Burckhardt I, Zimmermann S, Zeifang F, et al. Prosthetic infection: Improvement of diagnostic procedures using 16S ribosomal deoxyribonucleic acid polymerase chain reaction. *Int Orthop*. 2013;37:2515–2521. doi:10.1007/s00264-013-2038-7.
- [13] Dempsey KE, Riggio MP, Lennon A, Hannah VE, Ramage G, Allan D, et al. Identification of bacteria on the surface of clinically infected and non-infected prosthetic hip joints removed during revision arthroplasties by 16S rRNA gene sequencing and by microbiological culture. *Arthritis Res Ther*. 2007;9:R46. doi:10.1186/ar2201.
- [14] Xu Y, Rudkjøbing VB, Simonsen O, Pedersen C, Lorenzen J, Schönheyder HC, et al. Bacterial diversity in suspected prosthetic joint infections: an exploratory study using 16S rRNA gene analysis. *FEMS Immunol Med Microbiol*. 2012;65:291–304. doi:10.1111/j.1574-695X.2012.00949.x.
- [15] Omar M, Petri M, Hawi N, Krettke C, Eberhard J, Liodakis E. Higher sensitivity of swab polymerase chain reaction compared with tissue cultures for diagnosing periprosthetic joint infection. *J Orthop Surg*. 2018;26:230949901876529. doi:10.1177/2309499018765296.
- [16] Stoodley P, Conti SF, Demeo PJ, Nistico L, Melton-Kreft R, Johnson S, et al. Characterization of a mixed MRSA/MRSE biofilm in an explanted total ankle arthroplasty. *FEMS Immunol Med Microbiol*. 2011;62:66–74. doi:10.1111/j.1574-695X.2011.00793.x.
- [17] Ryu SY, Greenwood-Quaintance KE, Hanssen AD, Mandrekar JN, Patel R. Low sensitivity of periprosthetic tissue PCR for prosthetic knee infection diagnosis. *Diagn Microbiol Infect Dis*. 2014;79:448–453. doi:10.1016/j.diagmicrobio.2014.03.021.
- [18] Mariaux S, Tabin UF, Borens O. Diagnosis of persistent infection in prosthetic two-stage exchange: pcr analysis of sonication fluid from bone cement spacers. *J Bone Jt Infect*. 2017;2:218–223. doi:10.7150/jbji.23078.
- [19] Morgenstern C, Cabric S, Perka C, Trampuz A, Renz N. Synovial fluid multiplex PCR is superior to culture for detection of low-virulent pathogens causing periprosthetic joint infection. *Diagn Microbiol Infect Dis*. 2018;90:115–119. doi:10.1016/j.diagmicrobio.2017.10.016.
- [20] Gomez E, Cazanave C, Cunningham SA, Greenwood-Quaintance KE, Steckelberg JM, UhlJR, et al. Prosthetic joint infection diagnosis using broad-range PCR of biofilms dislodged from knee and hip arthroplasty surfaces using sonication. *J Clin Microbiol*. 2012;50:3501–3508. doi:10.1128/JCM.00834-12.
- [21] Jun Y, Jianghua L. Diagnosis of periprosthetic joint infection using polymerase chain reaction: an updated systematic review and meta-analysis. *Surg Infect (Larchmt)*. 2018;19:555–565. doi:10.1089/sur.2018.014.
- [22] Qu X, Zhai Z, Li H, Li H, Liu X, Zhu Z, et al. PCR-based diagnosis of prosthetic joint infection. *J Clin Microbiol*. 2013;51:2742–2746. doi:10.1128/JCM.00657-13.
- [23] Tarabichi M, Shohat N, Goswami K, Alvand A, Silibovsky R, Belden K, et al. Diagnosis of periprosthetic joint infection: the potential of next-generation sequencing. *J Bone Joint Surg Am*. 2018;100:147–154. doi:10.2106/JBJS.17.00434.
- [24] Tarabichi M, Shohat N, Goswami K, Parvizi J. Can next generation sequencing play a role in detecting pathogens in synovial fluid? *Bone Joint J*. 2018;100:127–133. doi:10.1302/0301-620X.100B2.BJJ-2017-0531.R2.
- [25] Haddad FS. Next generation sequencing: is this the moment? *Bone Joint J*. 2018;100-B:125–126. doi:10.1302/0301-620X.100B2.BJJ-2018-0057.
- [26] Tani S, Lepetsos P, Stylianakis A, Vlamis J, Birbas K, Kaklamanos I. Superiority of the sonication method against conventional periprosthetic tissue cultures for diagnosis of prosthetic joint infections. *Eur J Orthop Surg Traumatol*. 2018;28:51–57. doi:10.1007/s00590-017-2012-y.

- [27] Hischebeth GT, Gravius S, Buhr JK, Molitor E, Wimmer MD, Hoerauf A, et al. Novel diagnostics in revision arthroplasty: implant sonication and multiplex polymerase chain reaction. *J Vis Exp*. 2017. doi:10.3791/55147.
- [28] Rak M, Kavčič M, Trebše R, Čór A. Detection of bacteria with molecular methods in prosthetic joint infection: sonication fluid better than periprosthetic tissue. *Acta Orthop*. 2016;87:339–345. doi:10.3109/17453674.2016.1165558.
- [29] Huang Z, Wu Q, Fang X, Li W, Zhang C, Zeng H, et al. Comparison of culture and broad-range polymerase chain reaction methods for diagnosing periprosthetic joint infection: analysis of joint fluid, periprosthetic tissue, and sonicated fluid. *Int Orthop*. 2018;42:2035–2040. doi:10.1007/s00264-018-3827-9.
- [30] Liu K, Fu J, Yu B, Sun W, Chen J, Hao L. Meta-analysis of sonication prosthetic fluid PCR for diagnosing periprosthetic joint infection. *PLoS One*. 2018;13:e0196418. doi:10.1371/journal.pone.0196418.
- [31] Parvizi J, Gehrke T. Definition of periprosthetic joint infection. *J Arthroplasty*. 2014;29:1331. doi:10.1016/j.arth.2014.03.009.
- [32] Pupaibool J, Fulnecky EJ, Swords RL, Sistrunk WW, Haddow AD. Alpha-defensin—novel synovial fluid biomarker for the diagnosis of periprosthetic joint infection. *Int Orthop*. 2016;40:2447–2452. doi:10.1007/s00264-016-3306-0.
- [33] Li B, Chen F, Liu Y, Xu G. Synovial fluid α -defensin as a biomarker for peri-prosthetic joint infection: a systematic review and meta-analysis. *Surgical Infections*. 2017. doi:10.1089/sur.2017.006.
- [34] Pérez-Prieto D, Portillo ME, Puig-Verdié L, Alier A, Martínez S, Sorlí L, et al. C-reactive protein may misdiagnose prosthetic joint infections, particularly chronic and low-grade infections. *Int Orthop*. 2017;41:1315–1319. doi:10.1007/s00264-017-3430-5.



Authors: Óliver Marín-Peña, Majd Tarabichi, Karan Goswami

QUESTION 2: What is the preferred type of sample (tissue, fluid, etc.) for molecular analysis in the diagnosis of orthopaedic infections?

RECOMMENDATION: Several molecular methods have been developed in an effort to provide a viable culture-independent alternative for diagnosis of orthopaedic infections. However, due to the variation between studies with respect to the techniques and variety of samples collected, it remains difficult to recommend collection of one specimen type over another. While we cannot recommend a single molecular diagnostic test, careful assessment of the individual technique (location, volume, medium, temperature and transport) utilized is needed for appropriate collection and yield from the corresponding samples.

LEVEL OF EVIDENCE: Limited

DELEGATE VOTE: Agree: 87%, Disagree: 2%, Abstain: 11% (Super Majority, Strong Consensus)

RATIONALE

Identification of the infecting organism is imperative in the management of periprosthetic joint infection (PJI) [1,2]. Unfortunately, current methods, namely culture, have failed to perform at a level where the infecting organism is routinely identified, with up to half of PJIs yielding no known pathogen on microbiological culture [3–7]. Several molecular techniques have been examined to address this issue, however, no single technique has established itself to be superior to others. Furthermore, the optimal specimen type for maximizing the sensitivity and specificity of such technologies is an even greater dilemma.

Conventional cultures typically rely on synovial fluid from aspiration, when available, as well as multiple tissue samples obtained intraoperatively. Swabs have largely fallen out of favor with evidence demonstrating their lack of sensitivity and specificity [8]. Culture of sonicate fluid has shown some promise, however conflicting results and the need for specialized equipment preclude its routine use [9].

Synovial Fluid

Synovial fluid has been studied extensively as a source material for identifying the infective organism in PJI. When successfully obtained in the preoperative setting, it may provide the surgeon with crucial information to help guide further operative management of a patient with PJI. Various studies have reported on the performance of synovial fluid based molecular diagnostics in isolation or in parallel with other specimen types. In a study by Huan et al., samples of periprosthetic tissue, sonication fluid and synovial fluid were collected for both culture and 16S broad-range polymerase chain reaction (PCR). The authors concluded that PCR of sonication fluid and synovial fluid were significantly more sensitive than PCR of periprosthetic tissue alone, with no difference in specificity [10]. Multiple studies have shown superiority of synovial fluid PCR to conventional culture, however, these studies simply assessed synovial fluid with no direct comparison to other specimen types [4,11–13]. In contrast, a study comparing the combined sensitivity and specificity of joint fluid culture and serum C-reactive protein levels versus synovial fluid PCR demonstrated inferior results.

Periprosthetic Tissue

Periprosthetic tissue is a useful specimen due to its abundance, as opposed to synovial fluid which may only be present in limited quantities, if at all. A meta-analysis by Qu et al. comparing tissue, synovial fluid and sonication fluid concluded that tissue samples conferred the maximal sensitivity, while sonication fluid helped optimize specificity [14]. Other reports have claimed that tissue PCR is inferior to culture, however these studies focused on a comparison between sonicate fluid culture/PCR and tissue [15,16].

Swab

Swabs have been used in a limited fashion for molecular analysis. Omar et al. compared swabs sampled for 16S rRNA PCR with those sent for tissue culture, and showed a higher sensitivity in favor of swab PCR compared to culture. This is the only report assessing the utility of swabs for molecular diagnosis of PJI. However, no direct comparison was made to other specimen types in this study [17].

While 16S rRNA PCR forms the bulk of studies assessing the different specimen types, there are emerging reports of newer techniques such as next-generation sequencing that will also need to be further explored in order to delineate the optimal specimen type [18–20]. Emerging evidence suggests that the use of gauze or larger swabs that are able to potentially sample a greater intraoperative surface area may confer a better sequencing yield.

In conclusion, the optimal specimen type for molecular analysis of PJI remains unknown. There is significant heterogeneity between studies with regard to the techniques assessed as well as the samples analyzed. Careful assessment of specific techniques are advised when using these technologies as part of the diagnostic workup.

REFERENCES

- [1] Nodzo SR, Bauer T, Pottinger PS, Garrigues GE, Bedair H, Deirmengian CA, et al. Conventional diagnostic challenges in periprosthetic joint infection. *J Am Acad Orthop Surg*. 2015;23 Suppl:S18–S25. doi:10.5435/JAAOS-D-14-00385.
- [2] Parvizi J, Erkokcak OF, Della Valle CJ. Culture-negative periprosthetic joint infection. *J Bone Joint Surg Am*. 2014;96:430–436. doi:10.2106/JBJS.L.01793.
- [3] Baré J, MacDonald SJ, Bourne RB. Preoperative evaluations in revision total knee arthroplasty. *Clin Orthop Relat Res*. 2006;446:40–44. doi:10.1097/01.blo.0000218727.14097.d5.
- [4] Gallo J, Kolar M, Dendis M, Loveckova Y, Sauer P, Zapletalova J, et al. Culture and PCR analysis of joint fluid in the diagnosis of prosthetic joint infection. *New Microbiol*. 2008;31:97–104.
- [5] Gomez E, Cazanave C, Cunningham SA, Greenwood–Quaintance KE, Steckelberg JM, Uhl JR, et al. Prosthetic joint infection diagnosis using broad-range PCR of biofilms dislodged from knee and hip arthroplasty surfaces using sonication. *J Clin Microbiol*. 2012;50:3501–3508. doi:10.1128/JCM.00834–12.
- [6] Shanmugasundaram S, Ricciardi BF, Briggs TWR, Sussmann PS, Bostrom MP. Evaluation and management of periprosthetic joint infection—an international, multicenter study. *HSS J*. 2014;10:36–44. doi:10.1007/s11420-013-9366-4.
- [7] Spangehl MJ, Masri BA, O’Connell JX, Duncan CP. Prospective analysis of preoperative and intraoperative investigations for the diagnosis of infection at the sites of two hundred and two revision total hip arthroplasties. *J Bone Joint Surg Am*. 1999;81:672–683.
- [8] Aggarwal VK, Higuera C, Deirmengian G, Parvizi J, Austin MS. Swab cultures are not as effective as tissue cultures for diagnosis of periprosthetic joint infection. *Clin Orthop Relat Res*. 2013;471:3196–3203. doi:10.1007/s11999-013-2974-y.
- [9] Rothenberg AC, Wilson AE, Hayes JP, O’Malley MJ, Klatt BA. Sonication of arthroplasty implants improves accuracy of periprosthetic joint infection cultures. *Clin Orthop Relat Res*. 2017;475:1827–1836. doi:10.1007/s11999-017-5315-8.
- [10] Huang Z, Wu Q, Fang X, Li W, Zhang C, Zeng H, et al. Comparison of culture and broad-range polymerase chain reaction methods for diagnosing periprosthetic joint infection: analysis of joint fluid, periprosthetic tissue, and sonicated fluid. *Int Orthop*. 2018;42:2035–2040. doi:10.1007/s00264-018-3827-9.
- [11] Morgenstern C, Cabric S, Perka C, Trampuz A, Renz N. Synovial fluid multiplex PCR is superior to culture for detection of low-virulent pathogens causing periprosthetic joint infection. *Diagn Microbiol Infect Dis*. 2018;90:115–119. doi:10.1016/j.diagmicrobio.2017.10.016.
- [12] Janz V, Schoon J, Morgenstern C, Preininger B, Reinke S, Duda G, et al. Rapid detection of periprosthetic joint infection using a combination of 16s rDNA polymerase chain reaction and lateral flow immunoassay: a pilot study. *Bone Joint Res*. 2018;7:12–19. doi:10.1302/2046-3758.71.BJR-2017-0103.R2.
- [13] Lausmann C, Zahar A, Citak M, Brañes J, Schmidl S, Frommelt L, et al. Are there benefits in early diagnosis of prosthetic joint infection with multiplex polymerase chain reaction? *J Bone Jt Infect*. 2017;2:175–183. doi:10.7150/jbji.22062.
- [14] Qu X, Zhai Z, Li H, Li H, Liu X, Zhu Z, et al. PCR-based diagnosis of prosthetic joint infection. *J Clin Microbiol*. 2013;51:2742–2746. doi:10.1128/JCM.00657–13.
- [15] Rak M, Kavčič M, Trebše R, Čór A. Detection of bacteria with molecular methods in prosthetic joint infection: sonication fluid better than periprosthetic tissue. *Acta Orthop*. 2016;87:339–345. doi:10.3109/17453674.2016.1165558.
- [16] Ryu SY, Greenwood–Quaintance KE, Hanssen AD, Mandrekar JN, Patel R. Low sensitivity of periprosthetic tissue PCR for prosthetic knee infection diagnosis. *Diagn Microbiol Infect Dis*. 2014;79:448–453. doi:10.1016/j.diagmicrobio.2014.03.021.
- [17] Omar M, Petri M, Hawi N, Krettek C, Eberhard J, Liodakis E. Higher sensitivity of swab polymerase chain reaction compared with tissue cultures for diagnosing periprosthetic joint infection. *J Orthop Surg (Hong Kong)*. 2018;26:2309499018765296. doi:10.1177/2309499018765296.
- [18] Tarabichi M, Shohat N, Goswami K, Alvand A, Parvizi J. Diagnosis of periprosthetic joint infection: the potential of next-generation sequencing. *J Bone Joint Surg Am*. 2018;100:147–154.
- [19] Tarabichi M, Shohat N, Goswami K, Parvizi J. Can next generation sequencing play a role in detecting pathogens in synovial fluid? *Bone Joint J*. 2018;100-B:127–133. doi:10.1302/0301-620X.100B2.BJJ-2017-0531.R2.
- [20] Ivy MI, Thoendel MJ, Jeraldo PR, Greenwood–Quaintance KE, Hanssen AD, Abdel MP, et al. Direct detection and identification of prosthetic joint infection pathogens in synovial fluid by metagenomic shotgun sequencing. *J Clin Microbiol*. 2018;56. doi:10.1128/JCM.00402–18.



Authors: Pablo S. Corona, Daniel Monsalvo, Hamidreza Yazdi, Matias Vicente

QUESTION 3: What is the best diagnostic method for identifying a *C. acnes* surgical site infection/periprosthetic joint infection (SSI/PJI)?

RECOMMENDATION: Microbiological cultures, incubated for a prolonged period (up to 14 days) is currently regarded as the best diagnostic method for identifying *C. acnes*. Subculture in thioglycolate broth is believed to improve the yield of culture for *C. acnes*.

LEVEL OF EVIDENCE: Moderate

DELEGATE VOTE: Agree: 92%, Disagree: 3%, Abstain: 5% (Super Majority, Strong Consensus)

RATIONALE

C. acnes is a slow-growing, anaerobic, aerotolerant, non-sporulating, gram-positive bacillus [1]. It is part of the normal microbiome of the skin and resides in deeper layers [2]. The strains isolated in cases of invasive infections (especially in relation to orthopaedic implants) differ from those identified on the skin surface in their capacity to produce biofilms [3,4]. Diagnosing low-grade infection after total joint arthroplasty (TJA) is often highly complex, as clinical

symptomatology and diagnostic studies may conflict [5,6]. *C. acnes* is also a common contaminant of bacterial cultures, thus the significance of recovering this organism from periprosthetic specimens is not always clear [7].

Clinical Signs and Symptoms

Diagnosis of hip and knee PJI caused by *C. acnes* remains challenging. This is primarily due to its indolent nature, which results in pain and stiffness as major complaints, rather than in the more classic signs of infection [6–9].

Serum Biomarkers

Tebruegge et al. found that white blood cell (WBC) count was normal in 75% of orthopaedic *C. acnes* infections [10] and several studies indicate that serum erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP) have a low sensitivity in such low-grade infections [5,7,10–14]. In a study focused on *C. acnes* total knee arthroplasty (TKA) infections [8], Nodzo et al. found that ESR and CRP levels were statistically lower in the *C. acnes* PJI group, as compared to a *Staphylococcus aureus* TKA infections (ESR: 23 mm per hour vs. 56 mm per hour, CRP: 2.0 mg/dl vs. 5.9 mg/dl). In a prospective study by Grosso et al. [15] on 69 patients who underwent revision shoulder arthroplasty, serum IL-6 was not an effective marker for diagnosing infection.

Synovial Biomarkers

Synovial fluid leukocyte count and neutrophil percentage have been reported as having high sensitivity and specificity in diagnosing hip and knee PJI [16–18]. The utility of the proposed cutoff points in cases of low-grade infections is unknown [13,19]. In a recent study by Nodzo et al., comparing 16 TKAs due to *C. acnes* PJI to 30 *S. aureus* TKA infections [8], the authors found that the median synovial fluid WBC count in the *C. acnes* group was 19,950 cell/mm³. This was similar to the count in their *S. aureus* group (26,250 cell/mm³, $p = 0.31$), as was the median percentage of polymorphonuclear cells (PMNs) in the synovial fluid (95.5% vs. 95%, respectively, $p = 0.13$).

With regard to synovial IL-6, a recent investigation found a strong association between elevated synovial fluid IL-6 level and positive *C. acnes* culture [20] in cases of shoulder PJI.

The presence of leukocyte esterase (LE) in the synovial fluid has recently been proposed as a quick and effective marker for PJI [21]. Its utility in cases of low-grade infection has not been fully investigated. In a prospective study focused on shoulder arthroplasty, the sensitivity of LE was 30% and the specificity was 67%. *C. acnes* was isolated in 63% of all positive cultures.

Numerous studies posit alpha-defensin 1 (AD-1) as a valuable biomarker for diagnosis of PJI [22–25]. Although alpha-defensin has been proven useful regardless of organism type [26], its utility in cases of low-grade pathogens like *C. acnes* is a matter of debate. In a recent prospective study by Frangiamore et al., 33 cases of painful shoulder arthroplasty were evaluated for infection [27]. They found that alpha-defensin showed a sensitivity of 63%, a specificity of 95% and an area under the curve (AUC) of 0.78 for diagnosis of shoulder PJI. Although 63% sensitivity is not ideal for detecting all infections among infected cases, they found this an improvement over other preoperative tests. They also found a strong association between α -defensin levels and the growth of *C. acnes*, compared with a negative culture growth. The risk of having an α -defensin false-negative result [28] must be taken into account in such low-grade infections, along with the fact that the alpha-defensin test does not provide information on the identity of the infectious pathogen.

In summary, the utility of serum and synovial markers in the diagnosis of *C. acnes* periprosthetic joint infection remains unclear and in need of improvement.

Culture Techniques

C. acnes is a slow-growing, fastidious bacteria, which necessitates a longer incubation period than those routinely allowed for orthopaedic specimens. For a long time, *C. acnes* was underdiagnosed in bone and joint infections due to the short cultivation times routinely used in diagnostic laboratories [29–31]. In a study [8] comparing *C. acnes* TKA infections (16 cases) and *S. aureus* TKA infections (30 cases) the meantime for culture growth in the *C. acnes* group was 8.3 days, whereas it took a mean of 1.8 ± 0.8 days for $2.0 \pm S. aureus$ cultures to produce results ($p < 0.0001$). In another study, *C. acnes* cultures became positive at 3 to 27 days after surgery (45% of cultures were positive at 1 week, 86% at 2 weeks, 97% were positive at 3 weeks and 100% were positive at 4 weeks), so false-negative cultures for *C. acnes* may be as a result of short incubation or inadequate number of culture samples [11]. On the other hand, prolonging the incubation beyond a point (for instance beyond 14 days) may result in a high percentage of false-positive culture results, as *C. acnes* is a common contaminant of culture in microbiology laboratories.

It is common knowledge that *C. acnes* requires more than five incubation days to grow if routine cultures are used [32], but the best appropriate cultivation time is a point of controversy within the scientific community. Recent studies recommend a prolonged cultivation time – up to 14 days [31,33] – however, prolonging the incubation period is costly and labor-intensive and could also increase the likelihood of detecting organisms that are not clinically relevant. A recent study suggested that seven days of incubation should be sufficient for accurately diagnosing orthopaedic implant-associated infections [34]. In this study, 96.6% of the infections were detected within 7 days, however *C. acnes* caused only 1 out of the 58 infections studied. However, a study by Bossard et al. [30], focusing on 70 patients with *C. acnes* orthopaedic infections, found that reducing cultivation time to 7 days resulted in misdiagnosis in 15 patients (21.4%). Furthermore, the study showed that prolonging cultivation time beyond 10 days did not improve sensitivity. Thus, the authors recommend 10-day cultivation followed by a blind subculture in thioglycolate broth, in cases where suspicion of *C. acnes* infection is high. They found that thioglycolate broth culture of tissue biopsy specimens showed a significant difference in median time to positivity ($p = 0.0001$) as compared to other methods. Thioglycolate broth was most effective for the isolation *C. acnes* (sensitivity 66.3% in tissue samples and 75% in bone samples) with significantly different results than those for aerobic and anaerobic agar plates (sensitivity, 5.1% and 42.1%, respectively, $p = 0.0001$).

Culture for 10 days to isolate *C. acnes* is also supported by another study by Frangiamore et al. [35] evaluating shoulder arthroplasty patients. In a very recent study by Rieber et al., anaerobe culture became detectable in supplemented liver thioglycolate broth within six days, emphasizing the importance of using supplemented growth media to enhance detection of these pathogens [14].

There is a concern that longer incubation periods have the potential to yield false positive results due to specimen contamination, and may not be helpful for identifying true infections. In a study by Bossard et al., 61.7% of samples belonging to their no-infection group were recorded after day 7. These results are consistent with another study by Butler-Wu et al., which showed 21.7% of cases in which only 1 positive *C. acnes* sample labeled as no-

infection became positive after day 13 [31]. The proportion of positive cultures and the timing of culture growth may help to distinguish a true-positive from a false-positive result. In a retrospective study of 46 shoulder arthroplasty revision cases in which a positive *C. acnes* culture was identified, the time to culture growth was significantly shorter in the probable true-positive culture group ($p = 0.002$) compared with the probable contaminant group (median 5 days vs. 9 days). Significantly fewer days to culture growth were demonstrated among cases with a higher number of positive cultures ($p = 0.001$) and a higher proportion of positive cultures [35]. PJI specimens (true positives) were 6.3-times more likely to have 2 culture media positive for *C. acnes* growth than specimens from non-diagnostic events, and the authors considered a single culture-positive specimen in the absence of histologic findings to be non-diagnostic and most likely representing contamination [5,31].

Recent studies have suggested an improved effectiveness of the implant sonicate fluid culturing method over conventional periprosthetic tissue culture in detecting bacteria in total knee and total hip arthroplasty patients because of its ability to disrupt biofilm membranes [36]. Such superiority in cases of *C. acnes* infection is a matter of debate. A study conducted by Piper et al. [37], investigating the utility of implant sonication in 136 cases undergoing shoulder arthroplasty or resection, found that sonicate fluid culture was more sensitive than periprosthetic tissue culture for detection of definite prosthetic shoulder infection (66.7% vs. 54.5%, respectively, $p = 0.046$). A recent study by Portillo et al., investigating the sensitivity of sonication in 39 orthopaedic implant-associated infections – including 5 cases with *C. acnes* infection – detected all 5 *C. acnes* infections by sonication, but only 2 by conventional tissue cultures [38]. However, other authors have not found such advantages to the use of sonication in cases of *C. acnes* PJI. In a recent study by Bossard et al., which investigated the optimum cultivation time for isolation of *C. acnes* [30], sub-analysis of 35 cases with PJI caused by *C. acnes* found a 96.2% sensitivity for tissue biopsy specimens (25/26 cases) with at least 1 positive culture, as compared with sonication fluid at 46.2% (12/26). Grosso et al. evaluated the utility of implant sonication fluid cultures in diagnosing periprosthetic joint infection as compared with standard culture techniques in patients undergoing revision shoulder arthroplasty [39]. They found that implant sonication fluid cultures showed no significant superiority to standard intraoperative tissue and fluid cultures in the diagnosis of infection in patients undergoing revision shoulder arthroplasty.

Molecular Techniques

In recent years, several molecular tests that can detect the presence of pathogens by evaluating the genetic trace of these microorganisms have become available [40,41]. Such tests seem very promising, but they are also a target of ongoing criticism. One significant challenge for polymerase chain reaction (PCR) test is its inability to distinguish clinically important infections from mere traces of dead bacteria or bacteria that are part of the normal microbiota. Culture-independent techniques as species-specific PCR or broad-range 16S rDNA PCR have been used in the diagnosis of PJI. The high sensitivity in the detection of bacterial DNA and non-viable forms (useful in case of previous antimicrobial treatment) are described among its advantages [6,42,43]. In a recent study by Morgenstern et al., synovial fluid multiplex PCR was found superior to synovial fluid culture for detection of low-virulence bacteria such as *C. acnes* and coagulase-negative staphylococci [44]. Holmes et al. [41], developed a PCR-restriction fragment length polymorphism (RFLP) approach that identifies *C. acnes* in tissue specimens within a 24-hour period. This PCR-RFLP assay combines the sensitivity of PCR with the specificity of RFLP mapping to identify *C. acnes* in surgical isolates. The assay is robust and rapid and a *C. acnes*-positive tissue specimen can be confirmed within 24 hours of sampling, facilitating treatment decision making, targeted antibiotic therapy and monitoring to minimize implant failure and revision surgery [45].

However, they are not exempt from limitations. The limit of detection of the target sequence can be variable for each test, and in the absence of a quantitative technique, it can be difficult to determine whether a positive signal represents contamination or a clinically relevant infection. [6,42,43]. The universal PCR has difficulties in the case of polymicrobial infections and a low sensitivity for the diagnosis of PJI has been described [45,46].

The utility of molecular techniques, although promising, remains to be explored in the setting of *C. acnes* implant-associated infections [41,47]. Another new molecular technique that is gaining popularity is the use of next-generation sequencing (NGS) for identification of infecting pathogens causing PJI [48]. Based on a recent study from the Rothman Institute, NGS appeared to have a promising role in the identification of infecting organisms in over 80% of culture negative cases that included isolation of *C. acnes* in some cases. An ongoing study examining patients with shoulder pathophysiology at the same institution appears to indicate that NGS may be a better test than traditional culture for isolation of slow-growing organisms, such as *C. acnes* that result in PJI (data to be published soon).

Histologic Analysis

Frozen section histology of periprosthetic tissues has been recommended for patients undergoing revision hip or knee arthroplasty, for whom a diagnosis of PJI has not been established or has not been excluded [49]. There is a concern that low-virulence organisms like *C. acnes* could induce a less vigorous inflammatory reaction, characterized by a lower tissue concentration of neutrophils. According to data from a study by Grosso et al., frozen sections show a low sensitivity [50] in shoulder *C. acnes* infections (50%) using the diagnostic thresholds currently recommended for revision hip and knee arthroplasty (Feldman's criteria). The authors recommend a threshold of 10 polymorphonuclear leukocytes per 5 high-power fields, which results in an increased sensitivity (73%). In other instances, such as in a comparative study by Nodzo et al. [8], acute inflammation was identified in 88% of available tissue samples (14/16) in the TKA *C. acnes* infection group, as compared to 100% of samples (29/29) in the *S. aureus* group ($p = 0.05$).

REFERENCES

- [1] Gharamti AA, Kanafani ZA. Cutibacterium (formerly Propionibacterium) acnes infections associated with implantable devices. *Expert Rev Anti Infect Ther.* 2017;15:1083–1094. doi:10.1080/14787210.2017.1404452.
- [2] Achermann Y, Goldstein EJ, Coenye T, Shirtliff ME. Propionibacterium acnes: from commensal to opportunistic biofilm-associated implant pathogen. *Clin Microbiol Rev.* 2014;27:419–440. doi:10.1128/CMR.00092–13.
- [3] Perry A, Lambert P. Propionibacterium acnes: infection beyond the skin. *Expert Rev Anti Infect Ther.* 2011;9:1149–1156. doi:10.1586/eri.11.137.
- [4] Holmberg A, Lood R, Mörgelin M, Söderquist B, Holst E, Collin M, et al. Biofilm formation by Propionibacterium acnes is a characteristic of invasive isolates. *Clin Microbiol Infect.* 2009;15:787–795. doi:10.1111/j.1469-0691.2009.02747.x.
- [5] Figa R, Muñetón D, Gómez L, Matamala A, Lung M, Cuchi E, et al. Periprosthetic joint infection by Propionibacterium acnes: Clinical differences between monomicrobial versus polymicrobial infection. *Anaerobe.* 2017;44:143–149. doi:10.1016/j.anaerobe.2017.03.008.
- [6] Vasso M, Schiavone Panni A. Low-grade periprosthetic knee infection: diagnosis and management. *J Orthop Traumatol.* 2015;16:1–7. doi:10.1007/s10195-014-0294-y.
- [7] Lavergne V, Malo M, Gaudelli C, Laprade M, Leduc S, Laflamme P, et al. Clinical impact of positive Propionibacterium acnes cultures in orthopedic surgery. *Orthop Traumatol Surg Res.* 2017;103:307–314. doi:10.1016/j.otsr.2016.12.005.
- [8] Nodzo SR, Westrich GH, Henry MW, Miller AO. Clinical analysis of propionibacterium acnes infection after total knee arthroplasty. *J Arthroplasty.* 2016;31:1986–1989. doi:10.1016/j.arth.2016.02.025.
- [9] Shah NB, Tande AJ, Patel R, Barbari EF. Anaerobic prosthetic joint infection. *Anaerobe.* 2015;36:1–8. doi:10.1016/j.anaerobe.2015.08.003.

- [10] Tebruegge M, Jones C, de Graaf H, Sukhtankar P, Allan RN, Howlin RP, et al. Invasive propionibacterium acnes infections in a non-selective patient cohort: clinical manifestations, management and outcome. *Eur J Clin Microbiol Infect Dis*. 2015;34:527–534. doi:10.1007/s10096-014-2256-y.
- [11] Pottinger P, Butler-Wu S, Neradilek MB, Merritt A, Bertelsen A, Jette JL, et al. Prognostic factors for bacterial cultures positive for propionibacterium acnes and other organisms in a large series of revision shoulder arthroplasties performed for stiffness, pain, or loosening. *J Bone Joint Surg Am*. 2012;94:2075–2083. doi:10.2106/JBJS.K.00861.
- [12] Pérez-Prieto D, Portillo ME, Puig-Verdié L, Alier A, Martínez S, Sorlí L, et al. C-reactive protein may misdiagnose prosthetic joint infections, particularly chronic and low-grade infections. *Int Orthop*. 2017;41:1315–1319. doi:10.1007/s00264-017-3430-5.
- [13] McArthur BA, Abdel MP, Taunton MJ, Osmon DR, Hanssen AD. Seronegative infections in hip and knee arthroplasty: periprosthetic infections with normal erythrocyte sedimentation rate and C-reactive protein level. *Bone Joint J*. 2015;97-B:939–944. doi:10.1302/0301-620X.97B7.35500.
- [14] Rieber H, Frontzek A, Jerosch J, Alefeld M, Strohecker T, Ulatowski M, et al. Periprosthetic joint infection caused by anaerobes. Retrospective analysis reveals no need for prolonged cultivation time if sensitive supplemented growth media are used. *Anaerobe*. 2018;50:12–18. doi:10.1016/j.anaerobe.2018.01.009.
- [15] Grosso MJ, Frangiamore SJ, Saleh A, Kovac MF, Hayashi R, Ricchetti ET, et al. Poor utility of serum interleukin-6 levels to predict indolent periprosthetic shoulder infections. *J Shoulder Elbow Surg*. 2014;23:1277–1281. doi:10.1016/j.jse.2013.12.023.
- [16] Ghanem E, Parvizi J, Burnett RSJ, Sharkey PF, Keshavarzi N, Aggarwal A, et al. Cell count and differential of aspirated fluid in the diagnosis of infection at the site of total knee arthroplasty. *J Bone Joint Surg Am*. 2008;90:1637–1643. doi:10.2106/JBJS.G.00470.
- [17] Trampuz A, Hanssen AD, Osmon DR, Mandrekar J, Steckelberg JM, Patel R. Synovial fluid leukocyte count and differential for the diagnosis of prosthetic knee infection. *Am J Med*. 2004;117:556–562. doi:10.1016/j.amjmed.2004.06.022.
- [18] Dinneen A, Guyot A, Clements J, Bradley N. Synovial fluid white cell and differential count in the diagnosis or exclusion of prosthetic joint infection. *Bone Joint J*. 2013;95-B:554–557. doi:10.1302/0301-620X.95B4.30388.
- [19] Grau L, Gunder MA, Schneiderbauer M. Difficult-to-detect low-grade infections responsible for poor outcomes in total knee arthroplasty. *Am J Orthop*. 2017;46:E148–E153.
- [20] Frangiamore SJ, Saleh A, Kovac MF, Grosso MJ, Zhang X, Bauer TW, et al. Synovial fluid interleukin-6 as a predictor of periprosthetic shoulder infection. *J Bone Joint Surg Am*. 2015;97:63–70. doi:10.2106/JBJS.N.00104.
- [21] Parvizi J, Jacovides C, Antoci V, Ghanem E. Diagnosis of periprosthetic joint infection: the utility of a simple yet unappreciated enzyme. *J Bone Joint Surg Am*. 2011;93:2242–2248. doi:10.2106/JBJS.J.01413.
- [22] Bonanzinga T, Zahar A, Dütsch M, Lausmann C, Kendoff D, Gehrke T. How reliable is the alpha-defensin immunoassay test for diagnosing periprosthetic joint infection? a prospective study. *Clin Orthop Relat Res*. 2017;475:408–415. doi:10.1007/s11999-016-4906-0.
- [23] Gehrke T, Lausmann C, Citak M, Bonanzinga T, Frommelt L, Zahar A. The accuracy of the alpha defensin lateral flow device for diagnosis of periprosthetic joint infection: comparison with a gold standard. *J Bone Joint Surg Am*. 2018;100:42–48. doi:10.2106/JBJS.16.01522.
- [24] Bingham J, Clarke H, Spangehl M, Schwartz A, Beauchamp C, Goldberg B. The alpha defensin-1 biomarker assay can be used to evaluate the potentially infected total joint arthroplasty. *Clin Orthop Relat Res*. 2014;472:4006–4009. doi:10.1007/s11999-014-3900-7.
- [25] Frangiamore SJ, Gajewski ND, Saleh A, Farias-Kovac M, Barsoum WK, Higuera CA. α -Defensin accuracy to diagnose periprosthetic joint infection—best available test? *J Arthroplasty*. 2016;31:456–460. doi:10.1016/j.arth.2015.09.035.
- [26] Deirmengian C, Kardos K, Kilmartin P, Cameron A, Schiller K, Booth RE, et al. The alpha-defensin test for periprosthetic joint infection outperforms the leukocyte esterase test strip. *Clin Orthop Relat Res*. 2015;473:198–203. doi:10.1007/s11999-014-3722-7.
- [27] Frangiamore SJ, Saleh A, Grosso MJ, Kovac MF, Higuera CA, Iannotti JP, et al. α -Defensin as a predictor of periprosthetic shoulder infection. *J Shoulder Elbow Surg*. 2015;24:1021–1027. doi:10.1016/j.jse.2014.12.021.
- [28] Adams JR, Schwartz AJ. False-negative synovial alpha-defensin. *Arthroplast Today*. 2017;3:239–241. doi:10.1016/j.artd.2017.05.006.
- [29] Abdulmassih R, Makadia J, Como J, Paulson M, Min Z, Bhanot N. Propionibacterium acnes: time-to-positivity in standard bacterial culture from different anatomical sites. *J Clin Med Res*. 2016;8:916–918. doi:10.14740/jocmr2753w.
- [30] Bossard DA, Ledergerber B, Zingg PO, Gerber C, Zinkernagel AS, Zbinden R, et al. Optimal length of cultivation time for isolation of propionibacterium acnes in suspected bone and joint infections is more than 7 days. *J Clin Microbiol*. 2016;54:3043–3049. doi:10.1128/JCM.01435-16.
- [31] Butler-Wu SM, Burns EM, Pottinger PS, Magaret AS, Rakeman JL, Matsen FA, et al. Optimization of periprosthetic culture for diagnosis of Propionibacterium acnes prosthetic joint infection. *J Clin Microbiol*. 2011;49:2490–2495. doi:10.1128/JCM.00450-11.
- [32] Dodson CC, Craig EV, Cordasco FA, Dines DM, Dines JS, Dicarolo E, et al. Propionibacterium acnes infection after shoulder arthroplasty: a diagnostic challenge. *J Shoulder Elbow Surg*. 2010;19:303–307. doi:10.1016/j.jse.2009.07.065.
- [33] Schäfer P, Fink B, Sandow D, Margull A, Berger I, Frommelt L. Prolonged bacterial culture to identify late periprosthetic joint infection: a promising strategy. *Clin Infect Dis*. 2008;47:1403–1409. doi:10.1086/592973.
- [34] Schwotzer N, Wahl P, Fracheboud D, Gautier E, Chuard C. Optimal culture incubation time in orthopedic device-associated infections: a retrospective analysis of prolonged 14-day incubation. *J Clin Microbiol*. 2014;52:61–66. doi:10.1128/JCM.01766-13.
- [35] Frangiamore SJ, Saleh A, Grosso MJ, Alolabi B, Bauer TW, Iannotti JP, et al. Early versus late culture growth of Propionibacterium acnes in revision shoulder arthroplasty. *J Bone Joint Surg Am*. 2015;97:1149–1158. doi:10.2106/JBJS.N.00881.
- [36] Trampuz A, Piper KE, Jacobson MJ, Hanssen AD, Unni KK, Osmon DR, et al. Sonication of removed hip and knee prostheses for diagnosis of infection. *N Engl J Med*. 2007;357:654–663. doi:10.1056/NEJMoa061588.
- [37] Piper KE, Jacobson MJ, Cofield RH, Sperling JW, Sanchez-Sotelo J, Osmon DR, et al. Microbiologic diagnosis of prosthetic shoulder infection by use of implant sonication. *J Clin Microbiol*. 2009;47:1878–1884. doi:10.1128/JCM.01686-08.
- [38] Portillo ME, Salvadó M, Trampuz A, Siverio A, Alier A, Sorlí L, et al. Improved diagnosis of orthopedic implant-associated infection by inoculation of sonication fluid into blood culture bottles. *J Clin Microbiol*. 2015;53:1622–1627. doi:10.1128/JCM.03683-14.
- [39] Grosso MJ, Frangiamore SJ, Yakubek G, Bauer TW, Iannotti JP, Ricchetti ET. Performance of implant sonication culture for the diagnosis of periprosthetic shoulder infection. *J Shoulder Elbow Surg*. 2018;27:211–216. doi:10.1016/j.jse.2017.08.008.
- [40] Hartley JC, Harris KA. Molecular techniques for diagnosing prosthetic joint infections. *J Antimicrob Chemother*. 2014;69:i21–i24. doi:10.1093/jac/dku249.
- [41] Holmes S, Pena Diaz AM, Athwal GS, Faber KJ, O’Gorman DB. Neer Award 2017: a rapid method for detecting propionibacterium acnes in surgical biopsy specimens from the shoulder. *J Shoulder Elbow Surg*. 2017;26:179–185. doi:10.1016/j.jse.2016.10.001.
- [42] Drancourt M, Bollet C, Carlioz A, Martelin R, Gayral JP, Raoult D. 16S ribosomal DNA sequence analysis of a large collection of environmental and clinical unidentifiable bacterial isolates. *J Clin Microbiol*. 2000;38:3623–3630.
- [43] Peeters B, Herijgers P, Beuselink K, Peetermans WE, Herregods M-C, Desmet S, et al. Comparison of PCR-electrospray ionization mass spectrometry with 16S rRNA PCR and amplicon sequencing for detection of bacteria in excised heart valves. *J Clin Microbiol*. 2016;54:2825–2831. doi:10.1128/JCM.01240-16.
- [44] Morganstern C, Cabric S, Perka C, Trampuz A, Renz N. Synovial fluid multiplex PCR is superior to culture for detection of low-virulent pathogens causing periprosthetic joint infection. *Diagn Microbiol Infect Dis*. 2018;90:115–119. doi:10.1016/j.diagmicrobio.2017.10.016.
- [45] Bémer P, Plouzeau C, Tande D, Léger J, Giraudeau B, Valentin AS, et al. Evaluation of 16S rRNA gene PCR sensitivity and specificity for diagnosis of prosthetic joint infection: a prospective multicenter cross-sectional study. *J Clin Microbiol*. 2014;52:3583–589. doi:10.1128/JCM.01459-14.
- [46] Gomez E, Cazanave C, Cunningham SA, Greenwood-Quaintance KE, Steckelberg JM, Uhl JR, et al. Prosthetic joint infection diagnosis using broad-range PCR of biofilms dislodged from knee and hip arthroplasty surfaces using sonication. *J Clin Microbiol*. 2012;50:3501–3508. doi:10.1128/JCM.00834-12.
- [47] Tarabichi M, Shohat N, Goswami K, Parvizi J. Can next generation sequencing play a role in detecting pathogens in synovial fluid? *Bone Joint J*. 2018;100-B:127–133. doi:10.1302/0301-620X.100B2.BJJ-2017-0531.R2.
- [48] Tarabichi M, Shohat N, Goswami K, Alvand A, Silibovsky R, Belden K, et al. Diagnosis of periprosthetic joint infection: the potential of next-generation sequencing. *J Bone Joint Surg Am*. 2018;100:147–154. doi:10.2106/JBJS.17.00434.
- [49] Della Valle C, Parvizi J, Bauer TW, Dicesare PE, Evans RP, Segreti J, et al. Diagnosis of periprosthetic joint infections of the hip and knee. *J Am Acad Orthop Surg*. 2010;18:760–770.



Authors: Alexander Shope, Naomi Kobayashi, Adolfo Llinás

QUESTION 4: Should organisms (e.g., *Treponema spp.*, *Corynebacteria spp.*) identified through molecular or genetic testing be treated the same as the pathogens isolated by culture?

RECOMMENDATION: No. Because of their associated poor clinical outcomes, unusual organisms resulting in infection should not be treated equivalently to a usual pathogenic organism. Identification of unusual organisms through molecular and genetic techniques should help aid in antibiotic selection in conjunction with surgery, as indicated. Because of the associated poor clinical outcomes of unusual organisms and polymicrobial infections, the results of these newer techniques should not be ignored, but instead used to help inform therapeutic choices.

LEVEL OF EVIDENCE: Limited

DELEGATE VOTE: Agree: 93%, Disagree: 2%, Abstain: 5% (Super Majority, Strong Consensus)

RATIONALE

There are variety of unusual organisms that can cause periprosthetic joint infections (PJI) aside from *Staphylococcus* species. Unusual organisms represent about 4.5% of the PJIs in the United States, while culture-negative infections account for 18.6% [1]. Many of these uncommon organisms, in addition to the culture-negative organisms, are associated with polymicrobial PJIs [2]. In order to manage such patients, broad-spectrum antibiotics are often required that need tailored to the specific organisms causing the infection due to high rates of antibiotic resistance [2].

In recent a retrospective study, methicillin-resistant *Staphylococcus aureus* (MRSA), *Pseudomonas* and *Proteus*-related PJI have been associated with lower infection-free rates, which means more surgery and hospital time are required for definitive treatment [3]. Thus, aside from MRSA, there are other organisms that are associated with poor PJI outcomes.

In polymicrobial PJI, clinical outcomes were reported to be poor when compared to monomicrobial or culture-negative PJI [2]. In addition, polymicrobial PJI had higher rate of amputation (odds ratio (OR): 3.8, 95% confidence interval (CI) 1.34 to 10.80, $p = 0.012$), arthrodesis (OR: 11.06, 95% CI 1.27 to 96.00, $p = 0.029$) and PJI-related mortality (OR: 7.88, 95% CI 1.60 to 38.67, $p = 0.011$) compared with patients with monomicrobial PJI [2]. In such polymicrobial PJI, gram-negative organisms (OR: 6.33, $p < 0.01$), enterococci (OR: 11.36, $p < 0.01$), *Escherichia coli* (OR: 6.55, $p < 0.01$) and atypical organisms (OR: 9.85, $p < 0.01$) isolation were associated with polymicrobial PJIs [2]. PJI due to gram-negative species such as *Pseudomonas aeruginosa*, *Escherichia coli* and *Klebsiella pneumoniae* have proved to have lower rates of therapeutic success following debridement when compared to gram-positive organisms [4].

Fungal infection should also be recognized as an atypical organism causing PJI. Although the reports describing PJI due to fungal infection are limited, the clinical outcomes of PJI by *Candida* species were unsatisfactory. It was reported that the overall rate of mortality attributable to *Candida* PJI was 25% [5]. Multidrug-resistant gram-negative organisms, such as carbapenemase-producing *Klebsiella pneumoniae*, require aggressive medical and surgical treatment [6]. In a small case series of *Propionibacterium avidum* PJIs, debridement-retention of the prosthesis was not an effective option [7]. Similarly, although *Enterococcal* PJI is not frequent, its successful rate of treatment was reported to be low [8,9].

Because clinical outcomes can be associated with the characteristics of the causative agent, the ideal goal is to properly identify all pathogens responsible for the infection [2]. However, some of these unusual organisms can be difficult to detect or take excessive time to appropriately culture [10]. Negative culture results can pose a challenge for physicians therapeutically, for they lack vital diagnostic information, such as the true identity of the causative agent(s). Recently, research has focused on newer innovative methods of infection detection and identification. At the forefront of these new innovative techniques are molecular and genetic methods such as polymerase chain reaction (PCR) assay. Although current molecular and genetic methods tend to have high sensitivities, their specificities are lower and therefore cannot be used as a single diagnostic test as of now [10]. However, as technologies continue to improve, more insight into the pathologic agents will likely become available allowing physicians to make more informed therapeutic decisions based on information such as the presence of antibiotic resistant genes.

A study by Tarabichi et al. examined the utility of some of the newer molecular and genetic techniques, also known as next-generation sequencing (NGS) [11]. Based on the results of their study, they were able to conclude that NGS may be a useful adjunct to aid in organism identification [11]. Although their study shows much promise, they do note that further larger studies are needed to further validate this new technology.

Although two-stage exchange arthroplasty remains the gold standard for surgical management of chronic PJIs, especially when the causative organism is a resistant microbe or produces biofilm, the emergence of new pathogen identification methods will potentially allow physicians to choose more appropriate antibiotic regimens [9,11,12]. Much research is still needed for further validation of these techniques. However, it is clear that infection secondary to unusual organisms are associated with poor clinical outcomes and therefore should be treated with some variation from standard protocols, even if that is simply a more informed antibiotic regimen choice. Information from newer molecular and genetic techniques shows much promise in aiding in diagnosis of these types of infections.

REFERENCES

- [1] McLawhorn AS, Nawabi DH, Ranawat AS. Management of resistant, atypical and culture–negative periprosthetic joint infections after hip and knee arthroplasty. *Open Orthop J*. 2016;10:615–632. doi:10.2174/1874325001610010615.
- [2] Tan TL, Kheir MM, Tan DD, Parvizi J. Polymicrobial periprosthetic joint infections: outcome of treatment and identification of risk factors. *J Bone Joint Surg Am*. 2016;98:2082–8. doi:10.2106/JBJS.15.01450.
- [3] Cunningham DJ, Kavolus JJ, Bolognesi MP, Wellman SS, Seyler TM. Specific infectious organisms associated with poor outcomes in treatment for hip periprosthetic infection. *J Arthroplasty*. 2017;32:1984–1990.e5. doi:10.1016/j.arth.2017.01.027.
- [4] Hsieh P, Lee MS, Hsu K, Chang Y, Shih H, Ueng SW. Gram-negative prosthetic joint infections: risk factors and outcome of treatment. *Clin Infect Dis*. 2009;49:1036–1043. doi:10.1086/605593.
- [5] Ueng SW, Lee CY, Hu C, Hsieh PH, Chang Y. What is the success of treatment of hip and knee candidal periprosthetic joint infection? *Clin Orthop Relat Res*. 2013;471:3002–3009. doi:10.1007/s11999-013-3007-6.
- [6] de Sanctis J, Teixeira L, van Duin D, Odio C, Hall G, Tomford JW, et al. Complex prosthetic joint infections due to carbapenemase–producing *Klebsiella pneumoniae*: a unique challenge in the era of untreatable infections. *Int J Infect Dis*. 2014;25:73–78. doi:10.1016/j.ijid.2014.01.028.
- [7] Achermann Y, Liu J, Zbinden R, Zingg PO, Anagnostopoulos A, Barnard E, et al. *Propionibacterium avidum*: A virulent pathogen causing hip periprosthetic joint infection. *Clin Infect Dis*. 2018;66:54–63. doi:10.1093/cid/cix665.
- [8] Rasouli MR, Tripathi MS, Kenyon R, Wetters N, Della Valle CJ, Parvizi J. Low rate of infection control in enterococcal periprosthetic joint infections. *Clin Orthop Relat Res*. 2012;470:2708–2716. doi:10.1007/s11999-012-2374-8.
- [9] Kheir MM, Tan TL, Higuera C, George J, Della Valle CJ, Shen M, et al. Periprosthetic joint infections caused by enterococci have poor outcomes. *J Arthroplasty*. 2017;32:933–947. doi:10.1016/j.arth.2016.09.017.
- [10] Yoon HK, Cho SH, Lee DY, Kang BH, Lee SH, Moon DG, et al. A review of the literature on culture–negative periprosthetic joint infection: epidemiology, diagnosis and treatment. *Knee Surg Relat Res*. 2017;29:155–164. doi:10.5792/ksrr.16.034.
- [11] Tarabichi M, Shohat N, Goswami K, Alvand A, Silibovsky R, Belden K, et al. Diagnosis of periprosthetic joint infection: the potential of next–generation sequencing. *J Bone Joint Surg*. 2018;100:147–154. doi:10.2106/JBJS.17.00434.
- [12] Kapadia BH, Berg RA, Daley JA, Fritz J, Bhawe A, Mont MA. Periprosthetic joint infection. *Lancet*. 2016;387:386–394. doi:10.1016/S0140-6736(14)61798-0.