

## Perioperative Albumin in TKA: Age and Baseline as Dynamic Predictors

Yoshinori Ishii<sup>1</sup>, Hideo Noguchi<sup>1</sup>, Junko Sato<sup>1</sup>, Ikuko Takahashi<sup>1</sup>  
<sup>1</sup>Ishii Orthopaedic and Rehabilitation Clinic, Gyoda Saitama, JAPAN  
 Email; ishii@sakitama.or.jp

**Disclosures:** The authors declare that they have no competing interests

**INTRODUCTION:** Previous studies have evaluated preoperative serum albumin (SA) for predicting postoperative complications of total knee arthroplasty (TKA). This study aimed to investigate the dynamics of perioperative SA and changes in SA ( $\Delta$ SA) and identify any influential patient or surgical factors. **METHODS:** In total, 381 patients (483 knees) undergoing primary TKA were recruited (Table 1). SA values preoperatively ( $SA_0$ ), 1 week postoperatively ( $SA_{1W}$ ), and 4 weeks postoperatively ( $SA_{4W}$ ) were investigated. SA values were converted to a percentage of  $SA_0$  and differences between timepoints were calculated and expressed as follows:  $\Delta SA_{1W-0}$ ,  $\Delta SA_{4W-1W}$ , and  $\Delta SA_{4W-0}$ . Patient and surgical factors previously identified or with the potential to influence SA were evaluated.

**RESULTS:** The median values of  $SA_0$ ,  $SA_{1W}$ , and  $SA_{4W}$  were 4.4, 3.8, and 4.2 g/dL, respectively;  $SA_0$  was significantly different between groups ( $p < 0.001$ ) (Table 2). The incidence of low  $SA_0$  ( $< 3.5$  g/dL) was less than 1%. Median  $\Delta$ SA values were -13.7%, 9.6%, and -4.5% for  $\Delta SA_{1W-0}$ ,  $\Delta SA_{4W-1W}$  and  $\Delta SA_{4W-0}$ , respectively;  $\Delta$ SA was significantly different between groups ( $p < 0.001$ ) (Table 3).  $SA_{4W}$  recovered to 95.5% of  $SA_0$  with less than 2% of patients having low  $SA_{4W}$  ( $< 3.5$  g/dL). Multiple regression analyses showed SA concentration at each timepoint was significantly associated with the other SA timepoint values; age was significantly associated with  $SA_{4W}$  and  $SA_{1W}$  (all  $p < 0.001$ ).

**DISCUSSION:** We identified  $SA_0$  and age as significant factors affecting SA dynamics in the perioperative period. Low SA ( $< 3.5$  g/dL) was uncommon both preoperatively and at 4 weeks postoperatively; therefore, conventional cutoff values and preventive measures for low SA may need reconsideration.

**SIGNIFICANCE/CLINICAL RELEVANCE:** This study highlights that preoperative serum albumin and age are crucial factors influencing perioperative albumin dynamics after TKA, suggesting a need to reconsider conventional low albumin cutoff values and their associated preventive measures due to the rarity of severe postoperative drops.

**Table 1** Patient characteristics

	Variables (patients/ knees)	381/ 483
Patient Factor	Sex (Male 63/ 77, Female 318/ 406)	18 [22]/ 79 [88]
	Age (years)	74 [69, 78]
	Body weight (kg), Male/ Female	58 [53,66]
	Body height (cm)	151 [147, 155]
	Body mass index (kg/m <sup>2</sup> ), Male/ Female	26 [23, 28]
	HSS score	45 [37, 51]
	Comorbidity	
	Hypertension (Yes/No)	301/ 182
Hyperlipidemia (Yes/No)	95/ 388	
Diabetes Mellitus (Yes/No)	59/ 424	
Smoking history (Yes/No)	24/ 459	
Surgical Factor	Operative Time (min.)	55 [50, 62]
	Torniquet Time (min.)	58 [53, 65]
	Total blood loss (ml)	430 [200, 720]

Data are presented as n or median [25th, 75th percentile]. HSS, Hospital for Special Surgery.

**Table 2** Distribution of patients between serum albumin concentration ranges at each time point

Serum Albumin Level (g/dL)	Number of Patients (%)		
	Pre-op*	1 week post-op*	4 weeks post-op*
N (%)			
Median [quasi]	4.4 [4.2, 4.6]	3.8 [3.6, 3.9]	4.2 [4.0, 4.3]
< 3.0	2 (0.4)	3 (0.6)	1 (0.2)
3.0-3.4	2 (0.4)	60 (12.4)	8 (1.7)
3.5-3.9	38 (7.9)	301 (62.3)	93 (19.3)
4.0-4.4	251 (52.0)	117 (24.2)	310 (64.2)
$\geq 4.5$	190 (39.3)	2 (0.4)	71 (14.7)

\*  $p < 0.001$  (among three groups) using Friedman test, and each group ( $p < 0.001$ ) using Scheffé's method.

**Table 3** Distribution of patients between serum albumin (SA) concentration change ranges expressed as a percentage of preoperative SA

Period; N (%)	$\Delta$ Alb $\leq -20\%$	$-20 < \Delta$ Alb $\leq -10\%$	$-10 < \Delta$ Alb $\leq 0\%$	$0\% < \Delta$ Alb
(Po 1W- Pre)/ Pre -13.7 (-17.7, -9.5)*	58 (12.0)	291 (60.2)	126 (26.1)	8 (1.7)
(Po4w-Po1w)/ Pre 9.6 (5.7, 13.1)*	0 (0)	1 (0.2)	33 (6.8)	449 (93.0)
(Po 4w- Pre)/ Pre -4.5 (-8.2, -0.2)*	5 (1.0)	80 (16.6)	284 (58.8)	114 (23.6)

\*  $p < 0.001$  (among 3 groups) by the Friedman's test, and each group ( $p < 0.001$ ) with Scheffé's method